INDEX TO VOLUME XIX

New species in blackface type

ABBOTT, E. V., Diseases of economic plants in Peru, 645–656; Stem rust of wheat in Peru, 1041–1043

Abies, balsamea, butt-rot, 730; host of Pestalotia funerea, 203, 227

Abscission of young apples, relation to lime-sulphur sprays, 1001–1007

Abstracts of papers presented at the twentieth annual meeting of the American Phytopathological Society, New York, N. Y., December 28 to 31, 1928, inclusive, 79–108

Acidulated mercuric chloride, in disinfecting potato tubers for the control of Rhizoctonia, 713–724; loss of strength, 723; method, advantages, 722

Acrostalagmus aphidium, control of aphids by means of, 102

Actinomyces, cultural differences between species, including one undescribed, causing soil rot or pox of sweet potato, 183–187; p., new species causing pox of sweet potato, 183; undescribed species, cause of soil rot or pox of sweet potato, 179–190

Action, mass, in relation to infection with special reference to curly top of sugar beets, 1137

ADAMS, J. F., An Actinomycete the cause of soil rot or pox in sweet potatoes, 179–190

Adhesive plaster for control of root knots on grafted apple, 483–486

Adhesiveness of sulphur mixtures, 89

Aeciospores, germination of, 327–342

Aerial photographic methods, 1026

Agglutination tests with phytopathogenic bacteria, 99

Agronomic strains of timothy, reaction to Ustilago striaeformis, 105

Airplane photography in cotton root-rot study, 1025–1029

Alberta, Leptospaheria foot-rot of wheat in, 689–690

Alfalfa, black-stem disease of, 507–509; downy mildew, 651; leaf spot, 651; rust, 651; see Medicago sativa; white spot, 125

ALLARD, H. A., see GIDDINGS, N. J.

ALLEN, R. F., Heterothallism in Puccinia graminis, 1146–1147

Allium cepa, pink-root disease, 233–268; relation to H-ion concentration, 233–268

Alternaria, brassicae (Berk.) Sacc., leaf spot of cauliflower, 972; panax, control with calcium arsenate, 83; solani, 652; solani, physiological strains of, 533–548; tenuis, cotton leaf spot, 648

Amarillidaceae in California, leaf and scape spot of, 1138–1139

Amateur plant pathologist, 501

American Phytopathological Society, abstracts of papers presented at the twentieth annual meeting at New York, N. Y., December 28 to 31, 1928, inclusive, 79–108; minutes of field meeting of southern section, 531; report of the twentieth anniversary meeting, 511–526; second decade of, 503–506

Andes, plant-disease notes from, 969–974

Angular leaf spot of tobacco, 93

Anthocyan in infected corn leaves, 434

Anthracnose, of privet, 102; of raspberries, a new, 601–603; of snowberry, 101–102

Aphid, peach, as an agent in virus transmission, 109

Aphids, biologic control of, 102

Aphis gossypii, biologic control of, 102

Apical leaf roll of potato, 82–83

Aplanobacter michiganense, fruit spot tomato, 690

APOSTOLIDES, C. A., Leaf spot of sycamore caused by Stigmina platani (Fuckel) Sacc., 667–671

Apple, Armillaria mellea on, 869; diseases in Peru, 654; dropping, relation to lime-sulphur sprays, 1001–1007; fire blight, 86–87; fire-blight control and Bordeaux

spray, 285–293; fruit spot and surface rot, caused by Sporotrichum, 443–452; relation of fall applications of fungicides to scab control, 87–88; rust diseases of, 87; scab, control by dusts and sprays, 87; set and finish of fruits and blossom application of Bordeaux spray, 288; trees, control of root knots on grafted, 483–486; twig canker of, caused by Nectria cinnabarina, 1125–1128

Apricots, life history of Sclerotinia sclerotiorum in connection with the green rot of, 1136–1137

Arbor vitae, host of Pestalotia funerea, 203

Armillaria mellea, root rot of orchard trees, 869–873; studies on the nature of host resistance to, 1140–1141

Arsenate, calcium, as a fungicide, 83

Ascochyta, pisi, pinodella, lathyri, imperfecta, and phaseolorum on Leguminosae, 917–932

Ascochytae, host range and life history of some leguminous forms, 917–932

Aspergillus niger, carried by Leptoglossus zonatus, 481; cause of onion black mold, 733

Aster, wilt, control, 101; yellows, control, 101; yellows, control by wire-screen fences, 100; -yellows virus in insect vector, 1009–1015

Asters, wilt-resistant, 100–101

Attenuation of curly-top virus by resistant sugar beets which are symptomless carriers, 975–977

Avocado, certain diseases of, 1144; diseases in Peru, 654; tar spot, 972

Azalea, host of Pestalotia macrotricha, 214–215

BACH, W. J., see TAUBENHAUS, J. J.

Bacillus, amylovorus, adsorption by aluminium hydroxide, 291; amylovorus, control in apple and Bordeaux spray, 285–293; amylovorus, killing by copper dusts, 292–293; amylovorus, killing by lime water, 290; amylovorus, on apple, 86–87; atrosepticus, viability in soils, 295–300; carotovorus, viability in soils, 295–300

Bacteria, agglutination tests with, 99; differentiation of crown-gall and hairy-root types, 97–98; of crown-gall type, differentiation from nonpathogenic bacteria of radiobacter group, 98; relation to development of roots, 107

Bacterial, disease of corn, 81–82; leaf spot of sorghum, 82; parasites, influence of environment on, 96; parasites, influence of environment on infection by, 96; pustule of soy bean, 96; spot of peach, control of, 106–107; spot of radish and turnip, 97

Bacteriological differentiation of crown-gall and hairy-root types of bacteria, 97–98

Bacteriophage from Pseudomonas tumefaciens, 98

Bacterium, angulatum, on tobacco, 93; phaseoli, seed infection by, 96; phaseoli var. sojense on soy bean, 96; pruni, spray for control of, 106–107; tabacum, migration through leaf tissues, 97; translucens, physiological studies of, 99; translucens var. undulosum, physiological studies of, 99; tumefaciens, control on grafted apple trees, 483–486; tumefaciens, differentiation from nonpathogenic bacteria of radiobacter group, 98; tumefaciens, differentiation of crown-gall and hairy-root types, 97–98; tumefaciens, lytic principle from, 98; tumefaciens on hothouse roses, 1145; tumefaciens, relation of types of bacteria to root development, 107; tumefaciens, some researches on, 98–99

BAILEY, ALICE ALLEN, see RAMSEY, G. B.

BAILEY, D. L., President's address—retrospect and prospect, 413

BAIN, HENRY F., see SHEAR, C. L.; see STEVENS, NEIL E.

Balsam fir, butt-rot of, 725

Banana, anthracnose, 973; diseases in Peru, 654; host of Pestalotia leprogena, 217

BANFIELD, W. M., see RIKER, A. J.; see Wright, W. H.

Barley, control of covered smut on, 81; control of stripe disease, 81; Fusarium head blight of, 143; leaf scald, 651;

loose smut, 651; net blotch, 650; powdery mildew, 651; prevention of loose smut of, 413; response of hulless to seed treatment for covered smut and stripe disease, 657–666; scald, inheritance of resistance to, 1141; stem rust, 650; susceptibility of varieties to head blight, 80

BARRETT, J. T., Severe case of aerial "crown gall" on hothouse roses, 1145

BARSS, H. P., Some present-day problems, 493–501

Basal rot of narcissus bulbs, 99–100; relation to hot-water treatment, 100

Basisporium gallarum, corn-cob rot, 649

BEAL, J. A., see NELSON, RALPH M.

Bean, anthracnose, 973; diseases in Peru, 654; mosaic, virus of, longevity in seed, 605; seed infection by Bacterium phaseoli, 96

Beans, a Fusarium disease of, 84; effect of grafting on resistance and susceptibility to Colletotrichum lindemuthianum, 875–877

BEATTIE, R. KENT, biography of Nishida Tōji, 881–883

Beet, sugar, dry-rot canker of, 94–95; sugar, thinning as a means of controlling damping off, 95

Beetles, bark, relation of, to bluestain of southern pines, 1101–1106

BENNETT, C. W., Some raspberry mosaic symptoms, 89

Beta vulgaris, leaf condition with damping off, 317–318

Biologic control of aphids, 102

Biological abstracts, 494–495

BLACK, L. M., A new and striking abnormality in potato plants, 175–176

Black, locust, brooming disease of, 163; mold of onions, 733; mold of onions caused by Aspergillus niger v. Tiegh., 733–739; mold, onions, factors influencing infection, 736; shank of tobacco, control through disease resistance, 93; shank of tobacco in Porto Rico, 93–94; -stem disease of alfalfa, sweet clover, and red clover, 507–509

Blackfire of tobacco, 93

Blight, corn leaf, 649; early, potato, 652; head, susceptibility of barley varieties to, 80; late, potato, 651; leaf, coffee, 653; needle, of white pine, 413; seedling, influence of environment on predisposition of wheat and corn strains to, 79; seedling, inheritance of resistance to, in corn, 79–80; spur, of raspberries, 413

Blueberries, keeping quality, tests of, 593

Bluestain, fungi in southern pines, experiments with, 1101–1106; of wood, fungi causing, 597

Blue-staining fungi found in the United States, 597–599

Boll, rots, cotton, 648; spot, cotton, 648

BONDE, REINER, Physiological strains of Alternaria solani, 533–548; see SCHULTZ, E. S.

Book review, Principles of plant pathology, 177

Bordeaux, mixture, effect of, on the internal temperature of potato leaflets, 943–949; mixture, influence of form and proportion of lime and copper sulphate on suspension, 88; spray, effect as blossom spray on set and finish of apple fruits, 288; spray, in control of fire blight of apple, 285–293

Borer, European corn, 413

Botrytis, cultures of, 176

BOWMAN, JOHN J., see FULTON, HARRY R.

Bract spot, cotton, 648

Bremia lactucae, toxicity of commercial dusts to conidia of, 1143–1144

BRESSMAN, E. N., Effect of land plaster applied as a dust on seed corn, 1131–1133

BREWER, P. H., see KRAYBILL, H. R.; H. R. KRAYBILL, and M. W. GARDNER, Purification and certain properties of the tomato-mosaic virus, 108

Brooming disease of black locust (Robinia pseudacacia), 163–167

Brown spot, disease, pine needles, 993–999; pine needles, control of, 998

BRYAN, MARY K., Fruit spot of tomato caused by Aplanobacter michiganense, 690; The relation of moisture to stomatal infection, 751

Bud worm of balsam fir, predisposing factor for butt-rot infection, 731

Budding as a developmental phase common to many fungi, 1117–1123
Bulb, disease control of, 413
Bulbs, narcissus, relation of hot-water treatment to basal rot of, 100; narcissus, rot of, 99–100
Bunchy top of carnation, 973
BURGER, OWEN FRANCIS, In memoriam, 528
BURLISON, W. L., see HOLBERT, J. R.
BURRELL, A. B., see THOMAS, H. E.
Butt-rot of balsam fir, 725; caused by Polyporus balsameus Pk., 725–732; description, 727

Cabbage, ring spot of, 972
Cacoecia fumiferana, balsam fir, factor in butt-rot infection, 731
Caconema radicicola, see Heterodera radicicola
CAESAR, L., The European corn borer, 413
Calcium, arsenate, as a fungicide, 83; arsenate, as a fungicide for apple-scab control, 87–88; monosulphide, a substitute for lime sulphur for summer spraying, 106
CALDWELL, RALPH M., Preliminary results from cross inoculation and culture studies upon the fungus Rhynchosporium secalis (Oud.) Davis causing scald of cereals and other grasses, 104
Camellia, host of Pestalotia guepini, 199, 227; host of Pestalotia theae, 213
Cancer, chromosome number in tissues of, 97
Canker, dry-rot, of sugar beet, 94–95; twig, of apple caused by Nectria cinnabarina, 1125–1128
CANNON, W. N., see TISDALE, W. H.
Carapace spot of avocado, 1144
CARLETON, MARK ALFRED, biography of, 321–325
Carnation, bunchy top, 973
Carriers, symptomless, of sugar-beet curly-top virus, 975–977
CARSNER, EUBANKS, and C. F. LACKEY, Mass action in relation to infection with special reference to curly top of sugar beets, 1137
CARTER, WALTER, Ecological studies of curly top of sugar beets, 467–477
Carya, host of Pestalotia sphaerelloides, 207–227
CASH, LILLIAN, see JOHNSON, A. G.
Cauliflower, leaf spot of, 972
Cells, diseased, vacuolar changes in, 95
Cephalosporium leaf spot, coffee, 653
Ceratonia, host of Pestalotia curta, 210, 227
Ceratostomella pilifera, occurrence in U. S. A., 597
Cercospora, coffeicola, 653; coffeicola Berk. & Cke., 970; leaf spot of lettuce, 973; longissima (Cug.) Sacc., leaf spot on lettuce, 973; sorghi, corn leaf spot, 649; spot of coffee berry, 970
Cereal, scab development in 1928, 108
Cereals, infection of, with Ophiobolus graminis Sacc., 414; studies of the fungus causing scald of, 104
Chaetomella sacchari, sugar-cane-sheath disease, 647
Chamaecyparis, host of Pestalotia foedans, 206, 227; host of Pestalotia funerea, 203, 227
Chamaerops, host of Pestalotia palmarum, 212, 227
CHAMBERLIN, JOSEPH C., see KNIGHT, HUGH
CHARLES, VERA K., Coleodictyospora, a new genus of Dematiaceae, 1051
Chemical injury on watermelons, 85
Chemistry of the toxic factor of sulphur, 89
CHEN, H. K., see PORTER, R. H.
Cherry fruit, effect of sprays on weight of, 88
CHILDS, LEROY, and S. M. ZELLER, Observations on Armillaria root rot of orchard trees, 869–873
Chinese cucumbers, reaction to mosaic, 85–86
CHIVERS, A. H., A comparative study of Sclerotinia minor Jagger and Sclerotinia intermedia Ramsey in culture, 301–309
Chloroses of sugar cane, 647
CHRISTENSEN, J. J., The influence of temperature on the frequency of mutation in Helminthosporium sativum, 155–162; see STAKMAN, E. C.; H. A. RODENHISER, and CHIH TU, Susceptibility of barley

varieties to Fusarial head blight in Minnesota, 80
Chromosome number in crown-gall and cancer tissue, 97
Cicadula sexnotata, relation to aster-yellows virus, 1009
Citrullus, host of Pestalotia torulosa, 222, 227
Citrus, fruit diseases in Peru, 654; fruits, decay in transit, 89–90; further studies on Penicillia of, 1144; Nematospora on, 479–482; varieties and species of, relative resistance to bark diseases, 1136
Cladosporium fulvum, control by forced-air ventilation, 83; Cke., on tomato leaf, 973
CLAYTON, E. E., Breeding for resistance to cucumber-mosaic disease, 85; Toxicity of mercury and copper compounds in relation to their use for seed treatment and spraying, 86
Cliftonia, host of Pestalotia cliftoniae, 208, 227
CLINTON, G. P., Mosaic diseases of plants, 413; The new willow disease of Canada and New England, 413
Clover, black-stem disease, 507–509
Cob rot, corn, 649
Coconut, host of Pestalotia palmarum, 210–212
Coffee, brown spot, fruit and leaf, 653; Cercospora spot, 970; eye spot, 652; leaf blight, 653; leaf disease, 653; root disease, 653, 969; wilt, 653; zonal leaf spot, 653
Cold, chlorosis of sugar cane, 647; resistance and susceptibility in corn, 105–106
Coleodictyospora, new genus of Dematiaceae, 1051
College Station, Texas, report cotton root-rot conference at, 687–689
Colletotrichum, coffeanum, 653; falcatum, red rot sugar cane, 646; gloeosporioides Penz., on mango, 973; lindemuthianum, on beans, effect of grafting on resistance and susceptibility, 875–877; lindemuthianum (Sacc. and Magn.) Brio. & Cav., on bean pods, 973
Comptonia peregrina infected by rust, 453–454
Conference, cotton root-rot, report of, College Station, Texas, 687–689
Conifer, root and butt-rot, description, 746
Conifers, a fungous disease of, 91
Coniothyrium, phase of Leptosphaeria coniothyrium, 879
Conopholus americana, parasitism on oak roots, 102
Control, cranberry end rot, 1023; of bulb diseases, 413
Cocos, host of Pestalotia palmarum, 210, 227
COOK, MELVILLE T., The development of the spores of Plasmodiophora vascularum, 91–92
Cooperation, expansion of, 498–500
Copernica, host of Pestalotia copernica, 219, 227
Copper, compounds, toxicity in seed treatment and spraying, 86; dusts, killing of Bacillus amylovorus, 292–293; sulphate, influence of proportion of, on suspension of Bordeaux mixture, 88
Corn, cob rot, 649; disease of, 391–397; the European borer, 413; leaf blight, 649; leaf spot, 649; Penicillium injury, 105; preliminary report on bacterial disease of, 81–82; resistance and susceptibility to cold, 105–106; rust, 649; see maize; seed, effect of land plaster applied as a dust on, 1131–1133; seedling and root rots, 649; seedling blight, inheritance of resistance to, 79–80; seedling blight of, 79; smut, 649; smuts of, 415–442
Corypha, host of Pestalotia palmarum, 212, 227
Corticium, koleroga, 653; **C. praticola**, sp. nov., description of spore-forming Rhizoctonia, 1063–1065; **praticola**, sp. nov., effect of inorganic salts on the growth of, 1067–1068; **praticola**, sp. nov., effect of light on spore formation in, 1071; **praticola**, sp. nov., effect of moisture on spore formation in, 1069; **praticola**, sp. nov., effect of organic nitrogen on the growth of, 1068–1069; **praticola**, sp.

nov., effect of oxygen on spore formation, 1073–1074; **praticola**, sp. nov., effect of reaction of culture media on, 1071–1073; **praticola**, sp. nov., effect of temperature on the growth of, 1070; **praticola**, sp. nov., formal description of, 1065; **praticola**, sp. nov., growth and reproduction of, under controlled conditions, 1066–1067; **praticola**, sp. nov., heterothallism, 1074–1077, 1087–1088; **praticola**, sp. nov., isolation of single spores from individual basidia of, 1078–1084; **praticola**, sp. nov., mutations of, 1088–1090; **praticola**, sp. nov., parasitic on Medicago sativa, 1065; **praticola**, sp. nov., study of a new spore-forming Rhizoctonia, 1059–1099; **praticola**, sp. nov., sugars as food for, 1067; **praticola**, sp. nov., viability of basidiospores of, 1084–1087; C. vagum, cotton sore shin, 648; vagum, on sugar beet, 94–95

Cotton, boll rots, 648; boll spot, 648; bract spot, 648; growth disorders, 648; leaf spot, 648; mildew, 648; Nematospora on, 479–482; root knot, 649; root rot, method of inoculation for, 167; root rot, report conference, College Station, Texas, 687–689; root rot, study by airplane photography, 1025, 1029; sore shin, 648; Verticillium wilt of, 94; wilt, 647; wilt, a new, 171

COULSON, J. G., see MONTSERIN, B. G.

Covered smut, wheat, 650

Cowpea, susceptibility of varieties and selections to Fusarium wilt and root knot, 1145

Cranberry, end rot, 1017–1024; host of Pestalotia vaccinii, 201–202; relation of Godronia cassandrae Peck to, 1017; storage rots, 1928, 1037

Cronartium, comptoniae Arth., in western North America, 453–466; comptoniae, character of infection and damage, 454–462; comptoniae, distribution, 453–454; comptoniae, fruiting stages and spread, 462–463; comptoniae, hosts, 453–454; comptoniae, mycelium study, 742; comptoniae, western discovery, 453–454; filamentosum, aecia in the spring and fall, 462; ribicola, aecia in the spring and fall, 462; ribicola, on Pinus and Ribes in New York, 269–283

Crop, vegetable, diseases in Peru, 654–655

Crown gall, chromosome number in tissues of, 97; control of, 483–486; differentiation of crown-gall and hairy-root types of bacteria, 97–98; type of bacteria, differentiation from nonpathogenic forms, 98; viability of pathogene in soils, 295–300; "crown gall," aerial, on hothouse roses, 1145

Cruciferae, bacterial wilt of, 413

Cucumber, mosaic, breeding for resistance to, 85; mosaic virus, transmission by peach aphid, 109

Cucumbers, Chinese, reaction to mosaic, 85–86

Culture media, reaction of, effect on **Corticium praticola**, sp. nov., 1071–1073

Cultures, monosporidial, of Ustilago zeae, 420–434; of Ustilago zeae and Sorosporium reilianum, inoculation with mixed, 437–438

Cupressus, host of Pestalotia funerea, 203, 227

Curly top, attenuation of virus of, by resistant varieties, 975–977; development under artificial shade, 468–469; development under various light sources in the laboratory, 468–469; effect of the artificial control of environment on the development of, 471–477; of sugar beets, 1137; of sugar beets, ecological studies, 467–477; sugar beet, new host, 1031

Currant, death by white-pine blister rust a phase of control, 274

Cypress, host of Pestalotia foedans, 206; host of Pestalotia funerea, 203

Cytology of Ustilago zeae and Sorosporium reilianum, 91; studies in the physiology and, 415–442

Damage by Cronartium comptoniae, 454–462

Damping off, leaf condition caused in sugar beet, 317–318; thinning as a means of control in sugar beets, 95

DANA, B. F., see TAUBENHAUS, J. J.

Dasyscypha, agassizii (B. & C.) Sacc., occurrence, 579; agassizii, parasite of Pinus strobus, 277; arida Phillips, occurrence, 579; calyciformis, occurrence, 579; fusco-sanguinea, Pinus monticola affected by, 575, distribution, 578, morphology, 576; fusco-sanguinea Rehm, on western white pine, Pinus monticola Dougl., 575–584

Date palm, host of Pestalotia palmarum, 210

DAVIS, W. H., Anthracnose of snowberry (Symphoricarpos albus var. laevigatus), 101–102; Reaction in agronomic strains of timothy to Ustilago striaeformis (Westd.) Niessel, 105

DEAN, A. L., Root-observation boxes, 407–412

DEARNESS, J., Observations on needle blight of white pine, 413

Decay, of citrus fruits, 89–90; pines, caused by Lentinus lepideus, 710

Dematiaceae, a new genus of, 1051

Diamond canker of the prune, 1139

DICKSON, JAMES G., E. B. MAINS, and HELEN JOHANN, Progress report on cereal-scab development during the season of 1928, 108; P. E. HOPPE, J. R. HOLBERT, and GEORGE JANSSEN, The influence of environment during maturation upon predisposition to seedling blight in wheat and corn strains, 79; see LEUKEL, R. W.

DIEHL, W. W., see HASKELL, R. J.

Differentiation of crown-gall type of bacteria from nonpathogenic bacteria of radiobacter group, 98

Diplanetic species of Phytophthora, 92

Diplodia natalensis on citrus fruits in transit, 89–90

Diploid mycelium of Ustilago zeae, origin of, 430–431

Disease, -free seed and planting stock, 500; of conifers, related to snow cover, 91; of Lippia, caused by Sclerotium rolfsii Sacc., 509–510; of tobacco, caused by Septomyxa affinis, 90; phony, of peach, 107; resistance, control of black shank of tobacco through, 93; root, of pineapple and other plants due to **Tylenchus brachyurus**, n. sp., 611–629

Diseased cells, vacuolar changes in, 95

Diseases, bark, relative resistance of varieties and species of Citrus to, 1136; economic plants in Peru, 645–656; of sugar cane in Cuba, 343–366

Disinfectant, seed-grain, ethyl-mercury chloride as, 80

Disinfectants, liquid and dust for control of smuts on barley and wheat, 81

Dissociations and variability in the genus Fusarium, 753–868

DOAK, K. D., Parasitism of Conopholis americana Wallr. on roots of Quercus bicolor Willd., 102

DOBROSCKY, IRENE D., Is the aster-yellows virus detectable in its insect vector? 1009–1015

Douglas fir canker, 979

DRAYTON, F. L., Bulb growing in Holland in its relation to disease control, 413

DRECHSLER, CHARLES, A diplanetic species of Phytophthora causing pink rot of potato tubers, 92; A fruit rot of honeydew melons due to a species of Phytophthora, 85

Dry-rot canker of sugar beet, 94–95

DUFRÉNOY, J., Vacuolar changes in diseased cells, 95

DURRELL, L. W., Book review, Principles of plant pathology, 177

Dust disinfectants for control of smuts on barley and wheat, 81

Dusting, seed for oat smuts, 173; sulphur, prevention stem rust of wheat, 631–643

Dusts, commercial, toxicity to conidia of Bremia lactucae, 1143–1144; composed of lime sulphur and sulphur, 88–89; effect of, on seed corn, 1132; fungicidal action in relation to apple-scab control, 87

Dwarf, yellow, of onion, 86

Easter lily, mosaic, 311–315

Ecological studies of curly top of sugar beets, 467–477

EDDINS, A. H., Pathogenicity and cultural behavior of Ustilago zeae (Beckm.)

Ung., 885–916; Pathogenicity of multisporidial and monosporidial cultures of Ustilago zeae (Beckm.) Ung., 91

EDGECOMBE, A. E., see LINK, GEORGE K. K.

EDGERTON, C. W., E. C. TIMS, and P. J. MILLS, Relation of species of Pythium to the root-rot disease of sugar cane, 549–564

Effect, of Bordeaux mixture on the internal temperature of potato leaflets, 943–949; of grafting on resistance and susceptibility of beans to Colletotrichum lindemuthianum, 875–877; of Tilletia caries (DC.) Tul. (T. tritici (Bjerk.) Wint.) on development of wheat ear, 681–685

ELLIOTT, CHARLOTTE, Another bacterial leaf spot of sorghum, 82

End rot, cranberry, economic importance, 1022; of cranberry, 1017–1024

Environment, effect on the development of curly top, 471–477; influence on infection by bacterial parasites, 96

Environmental, factors, effect on root-knot nematode, 83–84; relations of **Tylenchus brachyurus**, n. sp., 626

Epidemiology of stem rust in relation to physiologic forms present, 951–959

Erianthus maximus var. Seemanni, leaf-splitting disease (Sclerospora northi West.), 961–967

Erwinia carotovora, following black mold, onions, 733

Erysiphe graminis, barley powdery mildew, 651

Ethyl-mercury chloride as a grain disinfectant, 80

Etiology, pink-root disease, onions, 691–704

Eucalyptus, host of Pestalotia inquinans, 225–227; host of Pestalotia molleriana, 223, 227

Eugenia, host of Pestalotia vermiformis, 224, 227

Eutettix, tenellus, ecological studies of, 467–477; tenellus, infected with curly top, filterable microorganism from, 1143; tenellus, relation to sugar-beet curly top, 1033

Exoascaceae, studies on, 90

Exoascus deformans, studies on, 90

Experimental error, method of equalizing in field plots, 565

Eye spot, coffee, 652; of sugar cane, 646

EZEKIEL, W. N., Report cotton-root-rot conference College Station, Texas, 687–689; see TAUBENHAUS, J. J.

Fall application of fungicides, relation to apple-scab control, 87–88

False smut of maize, Ustilaginoidea, 589–592

FAULL, J. H., A fungous disease of conifers related to the snow cover, 91

FAWCETT, HOWARD S., Nematospora on pomegranates, citrus, and cotton in California, 479–482; Occurrence and insect transmission of Nematospora in California, 1138; see KLOTZ, L. J.; and C. L. SHEAR, Discovery of the perfect stage of a Phomopsis on lemon bark, 1138

FELLOWS, HURLEY, Some chemical and morphological phenomena attending infection of the whole plant by Ophiobolus graminis Sacc., 103–104; Studies of certain soil phases of the wheat take-all problem, 103

Fellowships, graduate, 500

Fences, wire-screen, for control of aster yellows, 100

FERGUS, E. N., see VALLEAU, W. D.

Fiji, a new Sclerospora from, 961–967

Filterable microorganism from Eutettix tenellus and the sugar beet, both infected with curly top, 1143

Fir, Douglas, Armillaria mellea on, 869; host of Pestalotia funerea, 203

Fire blight, control in apple and Bordeaux spray, 285–293; notes on situation, 413; of apple, 86–87

Fomes pinicola, European and Canadian, 414; Fomes marginatus, same species as, 414; (Sw.), hosts of, 414

Foot rot, Leptosphaeria, of wheat in Alberta, 689–690

Forced-air ventilation for control of tomato-leaf mold, 83

Foreign-pest investigations, 497
FOSTER, W. R., see HENRY, A. W.
Freeing tobacco-mosaic virus from accompanying solids, 107–108
FREEMAN, E. M., Biography of Mark Alfred Carleton, 321–325
Fruit, citrus, decay in transit, 89–90; spot and surface rot of apple, caused by Sporotrichum, 443–452; spot, brown, coffee, 653; spot of tomato, caused by Aplanobacter michiganense, 690
Fruiting stages of Cronartium comptoniae, 462–464
Fruits, small, keeping quality affected by temperature, 593
FULTON, HARRY R., and JOHN J. BOWMAN, Decay of citrus fruits in transit, 89–90
Fungi, bluestain, in southern pines, 1101–1106; blue-staining, in U. S. A., 597; certain developmental phases common to many, 1117–1123
Fungicide, calcium arsenate as, 83
Fungicides, fall application in relation to apple-scab control, 87–88; research on, 497–498
Further studies on the modification of sugar-beet curly-top virus by its various hosts, 1141–1142
Fusarial head blight, susceptibility of varieties of barley to, 80
Fusarium, avenaceum, physiologic specialization in, 145; batatatis, control of, 84; cromyopthoron, factor in pink root of onions, 696; cromyopthoron, relation to H-ion concentration, temperature, and other cultural conditions, and to pink-root disease of onion, 233–268; culmorum, physiologic specialization in, 145; disease of beans, 84; graminearum, physiologic specialization in, 145; malli, factor in pink root of onions, 696; nivale, physiologic specialization in, 147; oxysporum, 652; pineapple root rots, caused by species of, 1146; rhizochromatistes, factor in pink root of onions, 696; (76 species and many strains), studies on variability and dissociation in, 753–868; solani, physiologic specialization in, 147; sp., basal rot of narcissus bulbs, caused by, 99–100; spp., physiologic specialization in, 143; vasinfectum, cotton wilt, 647; vasinfectum, in cotton, 171; wilt, coffee, 653; wilt, susceptibility of cowpea varieties and selections to, 1145
Fusicoccum putrefaciens Shear, relation to Godronia cassandrae Peck, 1017

GARDNER, M. W., see BREWER, P. H.; see KRAYBILL, H. R.; see WHITE, HAROLD E.; Sporotrichum fruit spot and surface rot of apple, 443–452
GARDNER, W. A., see JOHNSON, A. G.
Gaultheria, host of Pestalotia gibbosa, 216, 227
Germination, spores of Ustilago zeae, 419–420
Gibberella saubinetii, inheritance of resistance to, 79–80; physiologic specialization in, 143
GIDDINGS, N. J., H. A. ALLARD, and B. H. HITE, Inactivation of the tobacco-mosaic virus by high pressures, 749–750
GILBERT, A. H., Net necrosis of Irish potato tubers, 82
GILBERT, W. W., and F. C. MEIER, Chemical injury to watermelon in transit—results of shipping tests, 85
Gloeosporium, allantosporum Fautrey, raspberry anthracnose due to, 601; musarum Cke. & Massee on banana, 973; mutation in, 605
Glomerella cingulata on privet, 102
GODFREY, G. H., A nematode root lesion of pineapple and other crops, 94; Destructive root disease of pineapple and other plants due to **Tylenchus brachyurus**, n. sp., 611–629; and HELENE T. MORITA, Effects of some environmental factors on the root-knot nematode, 83–84
GODKIN, JAMES G., Physiological studies of Bacterium translucens and Bact. translucens var. undulosum, 99; see LINK, GEORGE K. K.
Godronia cassandrae Peck on cranberry, 1017–1024
Graduate fellowships, 500
Grafting, effect on resistance and susceptibility of beans to Colletotrichum lindemuthianum, 875–877

Grains, small, Fusarium head blight of, 143
Grape, host of Pestalotia menezesiana, 219; host of Pestalotia pezizoides, 197; wound gum of, 413
Graphium rigidum, occurrence in U. S. A., 597
Green rot of apricots, life history of Sclerotinia sclerotiorum in connection with, 1136–1137
Growth disorders, cotton, 648
GUBA, E. F., Monograph of the genus Pestalotia de Notaris, Part I, 191–232
Gum tree, host of Pestalotia inquinans, 225; tree, host of Pestalotia molleriana, 223; wound, due to fungi, 413
GUTERMAN, CARL E. F., see OGILVIE, LAWRENCE
Gymnosporangium, germinale, on apple, 87; globosum, on apple, 87; juniperi-virginianae, on apple, 87

HAASIS, FERDINAND W., see HARTLEY, CARL
Hadromycosis, due to Verticillium spp., world's literature on, 1138
HAHN, GLENN GARDNER, see WILSON, MALCOLM
Hairy root, control of, 483–486; differentiation of crown-gall and hairy-root types of bacteria, 97–98; relation of bacteria to development of, 107
HAMILTON, J. M., Studies of the fungicidal action of certain dusts and sprays in the control of apple scab, 87
HANNA, W. F., see STAKMAN, E. C.; Studies in the physiology and cytology of Ustilago zeae and Sorosporium reilianum, 91, 413, 415–442
HANSEN, H. N., Etiology of pink-root disease of onions, 691–704; **Phoma terrestris**, n. sp., as the cause of pink root of onions, 1135–1136
HARTER, L. L., A Fusarium disease of beans, 84; and W. A. WHITNEY, Masking of sweet-potato mosaic, 933–942
HARTLEY, CARL, Plot arrangement to distribute neighbor influence in field trials, 565–574; and FERDINAND W. HAASIS, Brooming disease of black locust (Robinia pseudacacia), 163–167
HASKELL, R. J., Report of the twentieth anniversary meeting of the American Phytopathological Society, 511–526; The second decade of the American Phytopathological Society, 503–506; and W. W. DIEHL, False smut of maize, Ustilaginoidea, 589–592
Head blight, Fusarium, of small grains, 143
Heart rot, living pines, caused by Lentinus lepideus, 705
HEDGCOCK, GEORGE G., Septoria acicola and the brown-spot disease of pine needles, 993–999
Helminthosporium, gossypii, leaf, bract, and boll spot of cotton, 648; ocellum, eye spot of sugar cane, 646; sativum, influence of temperature on the frequency of mutation in, 155–162; teres, net blotch of barley, 650; turcicum, corn leaf blight, 649
HENDERSON, W. J., see MELHUS, I. E.
HENRY, A. W., and W. R. FOSTER, Leptosphaeria foot rot of wheat in Alberta, 689–690
Heterodera radicicola, cotton root knot, 649; radicicola, effects of some environmental factors on, 83–84
Heterosporium echinulatum (Berk.) Cke., on carnation, 973
Heterothallic, forms of Fomes pinicola shown to be, 414
Heterothallism, in **Corticium praticola**, sp. nov., 1074–1077, 1087–1088; in Puccinia graminis, 1146–1147; in Sorosporium reilianum, 436–437; in Ustilago zeae, 885–916
HIGGINS, B. B., and V. H. YOUNG, Minutes of the field meeting of the southern section of the American Phytopathological Society, 531
High pressure, inactivating tobacco-mosaic virus, 749
HILL, J. BEN, Migration of Bacterium tabacum through the leaf tissues of Nicotiana tabacum, 97
History and distribution, Phomopsis pseudotsugae in Europe, 979–992

HITE, B. H., see GIDDINGS, N. J.
HOGGAN, ISMÉ A., The peach aphid (Myzus persicae Sulz.) as an agent in virus transmission, 109–123
HOLBERT, J. R., see DICKSON, JAMES G.; and W. L. BURLISON, Studies of cold resistance and susceptibility in corn, 105–106
Holland, bulb growing in, 413
Holly, host of Pestalotia stellata, 220
HOLMES, F. O., Local symptoms of mosaic in the leaves of some Nicotiana species, 92–93
HOPPE, P. E., Inheritance of resistance to seedling blight of corn caused by Gibberella saubinetii, 79–80; see DICKSON, JAMES G.
HORNE, WM. T., Progress in the study of certain diseases of the avocado, 1144
HORSFALL, JAMES G., Dusting seed for oat smuts, 173
Host range and life-history studies of some leguminous Ascochytae, 917–932
Hosts, distribution, Lentinus lepideus, 711
Hot-water treatment, relation to basal rot of narcissus bulbs, 100
Howea, host of Pestalotia palmarum, 210, 227
HOWITT, J. E., and R. E. STONE, A preliminary report of the results of some field tests in the use of seed disinfectants for the prevention of loose smut of barley, 413
HOWLETT, F. S., and CURTIS MAY, The relation of lime-sulphur sprays to the abscission of young apples, 1001–1007
HUBERT, ERNEST E., A butt-rot of balsam fir caused by Polyporus balsamaeus Pk., 725–732; A root and butt-rot of conifers caused by Polyporus circinatus Fr., 745–747
Humidity, relation to loose smut of wheat, 103
HUNGERFORD, C. W., see PIERCE, W. H.
HURT, R. H., Calcium monosulphide, a substitute for lime sulphur for summer spraying, 106
HUTCHINS, LEE M., Phony disease of the peach, 107
HUTCHINSON, W. G., Studies on the mycelium of Cronartium comptoniae Arthur on Pinus sylvestris L., 741–744
Hydrogen-ion concentration studies on Penicillium italicum and P. digitatum, 1142

Ilex, host of Pestalotia stellata, 220, 227
In memoriam, 527–529
Inactivation of the tobacco-mosaic virus by high pressures, 749–750
Incubation period for repeating pine rusts, 332
Infection, by Cronartium comptoniae, 454–462; by Ophiobolus graminis, 103–104; by Ustilago zeae of uninjured plants, 428–429; seed, by Bacterium phaseoli, 96
Inheritance of resistance to barley scald, 1141
Injury to corn due to Penicillium oxalicum, 105
Inoculation, a method of, for Phymatotrichum root-rot investigations, 167–170; tests with Septoria acicola, 994; with mixed cultures of Ustilago zeae and Sorosporium reilianum, 437–438
International conference for phytopathology and economic entomology—Holland, 1929, 607
Iodine, dust for oat-smut control, 173
Ipomoea batatas, masking of mosaic symptoms on, 933–942; soil rot or pox caused by Actinomyces, undescribed species, 179–190
Iris rust, 101
Irrigation, relation to white spot of alfalfa, 125–141

JANSSEN, GEORGE, see DICKSON, JAMES G.
JOHANN, HELEN, Further studies on Penicillium injury to corn, 105; see DICKSON, JAMES G.
JOHNSON, A. G., see LEUKEL, R. W.; LILLIAN CASH, and W. A. GARDNER, Preliminary report on a bacterial disease of corn, 81–82
JOHNSON, H. W., see LEACH, J. G.
JONES, D. H., A severe outbreak of bacterial wilt of Cruciferae, 413; Notes on the fire-blight situation, 413

JONES, L. R., and REGINA S. RIKER, Progress with the control of aster wilt and yellows, 101

JONES, WALTER, and T. E. RAWLINS, Influence of spindle-tuber disease on the physiology of the potato tuber, 1137

Keeping quality, Howes and McFarlin cranberries, 1039

KEITT, G. W., see RIKER, A. J.; see WILSON, E. E.; and E. E. WILSON, Third progress report of studies of fall applications of fungicides in relation to apple-scab control, 87–88

KENDRICK, JAMES B., Susceptibility of cowpea varieties and selections to Fusarium wilt and root knot, 1145; Toxicity of some commercial dusts to conidia of Bremia lactucae, 1143–1144

Kentia, host of Pestalotia palmarum, 210, 212, 227

KIRBY, R. S., Relation of the removal of smut balls to the control of stinking smut in winter wheat, 79

KLOTZ, L. J., Further studies on Penicillia of Citrus, 1144; and H. S. FAWCETT, Relative resistance of varieties and species of Citrus to bark diseases, 1136

KNIGHT, HUGH, JOSEPH C. CHAMBERLIN, and CHAS. D. SAMUELS, On some limiting factors in the use of saturated petroleum oils as insecticides on living plants, 1136

KOCH, L. W., The spur blight of raspberries, 413

KOTILA, JOHN E., Study of the biology of a new spore-forming Rhizoctonia, **Corticium praticola**, n. sp., 1059–1099

KRAYBILL, H. R., see BREWER, P. H.; P. H. BREWER, R. W. SAMSON, and M. W. GARDNER, The separation from mosaic tomato plants of the toxins which produce some of the typical mosaic symptoms, 108

KUNKEL, L. O., Wilt-resistant asters, 100–101; Wire-screen fences for the control of aster yellows, 100

LACHMUND, H. G., Cronartium comptoniae Arth. in western North America, 453–466

LACKEY, C. F., Attenuation of curly-top virus by resistant sugar beets, which are symptomless carriers, 975–977; Further studies of the modification of sugar-beet curly-top virus by its various hosts, 1141–1142; see CARSNER, EUBANKS

LAMBERT, EDMUND B., The relation of weather to the development of stem rust in the Mississippi Valley, 1–71; and E. C. STAKMAN, Sulphur dusting for prevention stem rust wheat, 631–643

Land plaster, effect of, applied as dust on seed corn, 1131–1133

Laurel, host of Pestalotia macrotricha, 214–215; host of Pestalotia rhododendri, 216

LEACH, J. G., Effect of grafting on resistance and susceptibility of beans to Colletotrichum lindemuthianum, 875–877; H. W. JOHNSON, and H. E. PARSONS, The use of acidulated mercuric chloride in disinfecting potato tubers for control of Rhizoctonia, 713–724

LEACH, LYSLE D., Observations on powdery mildew of raspberry in California, 1144–1145

Leaf, and scape spot of the Amarillidaceae in California, 1138–1139; blight, corn, 649; disease of coffee, 653; hopper, insect vector of aster-yellows virus, 1009–1015; mold of tomato, 973; mold of tomato, control by forced-air ventilation, 83; roll, apical, of potato, 82–83; rust, wheat, 650; scald, barley, 651; spot, alfalfa, 651; spot, bacterial, of sorghum, 82; spot, coffee, 653; spot, corn, 649; spot, cotton, 648; spot of cauliflower, 972; spot of Sycamore caused by Stigmina platani, 667–671; spot of tomato, 92; spot, zonal, coffee, 653; tissues, migration of Bacterium tabacum through, 97

Leaves, measuring surface of, 399–405

Leguminosae, Ascochytae on, 917–932

LEHMAN, S. G., Studies on bacterial pustule of soy bean, 96

Lemon bark, discovery of the perfect stage of a Phomopsis on, 1138

Lentinus lepideus, cause, heart rot, pines, 705–712; distribution and hosts, 711

LEONIAN, LEON H., Studies on the variability and dissociation in the genus Fusarium, 753–868
Leptoglossus zonatus carries fungi, 479–481
Leptosphaeria, coniothyrium, polymorphism of, 879; foot rot of wheat in Alberta, 689–690; sacchari, sugar-cane ring spot, 646
Lettuce, leaf spot of, 973
LEUKEL, R. W., Experiments with liquid and dust disinfectants for controlling covered smut of barley and stinking smut of wheat, 1926–1928, 81; J. G. DICKSON, and A. G. JOHNSON, Experiments on stripe disease of barley and its control, 81
LEVINE, M. N., see STAKMAN, E. C.
LEVINE, MICHAEL, The chromosome number in crown-gall and cancer tissues, 97
Life history, of Godronia cassandrae Peck, 1017; of Sphaceloma ampelinum, 673–679
Light, effect of, on spore formation in **Corticium praticola**, sp. nov., 1071
Lilium harrisii, mosaic, 311–315
Lime, influence of proportion of, on suspension of Bordeaux mixture, 88; sulphur and sulphur, dusts composed of, 88–89; sulphur, calcium monosulphides as a substitute for, 106; sulphur sprays, relation to abscission of young apples, 1001–1007
LINK, GEORGE K. K., A. E. EDGECOMBE, and J. G. GODKIN, Further agglutination tests with phytopathogenic bacteria, 99
Lippia canescens, a sclerotial disease of, 509–510
Liquid disinfectants for control of smuts on barley and wheat, 81
Literature, world's on hadromycosis due to Verticillium spp., 1138
Lobelia erinus, Rhizoctonia infecting, 585
Local symptoms of mosaic on species of Nicotiana, 92–93
Locust, black, brooming disease of, 163
LOH, T. C., see WHETZEL, H. H.
Longevity of bean-mosaic virus, 605
Loose smut, barley, 413, 651; wheat, 650
LUSK, J. P., see TAUBENHAUS, J. J.
Lyman memorial fund, 608; financial statement September, 1929, 880
Lytic principles from Pseudomonas tumefaciens, 98

MACHACEK, J. E., Black mold of onions, 413; The black mold of onions, caused by Aspergillus niger v. Tiegh., 733–739
MACKIE, W. W., Inheritance of resistance to barley scald, 1141; Resistance to Septoria tritici in wheat, 1139–1140
MAINS, E. B., Iris rust, Puccinia iridis, 101; Relative susceptibility of various varieties of sorghum to rust, Puccinia purpurea, 104; see DICKSON, JAMES G.
Maize, disease of, 391–397; Ustilaginoidea causing false smut, 589
Mangifera, host of Pestalotia virgatula, 222, 227
Mango, host of Pestalotia virgatula, 222; wither tip of, 973
Market pathology and transportation, 500
MARLOTH, RAIMUND H., Hydrogen-ion-concentration studies on Penicillium italicum and P. digitatum, 1142
MARTIN, G. HAMILTON, Certain developmental phases common to many fungi, 1117–1123; Polymorphism of Leptosphaeria coniothyrium (Fckl.) Sacc., 879
Masking of sweet-potato mosaic, 933–942
MAY, CURTIS, see HOWLETT, F. S.; see TILFORD, PAUL E., 943
MCCALLAN, S. E. A., see WHETZEL, H. H.
MCCALLUM, A. W., Woodgate rust in Canada, 414
MCCLUNG, C. E., Handbook of microscopical technique, review of, 609–610
MCCOWN, MONROE, Bordeaux spray in the control of fire blight of apple, 285–293
MCCUBBIN, W. A., Plant quarantines and the State, 487–492
Media, acid, sterilization of liquid, 413
Medicago sativa, **Corticium praticola**, sp. nov., parasitic on, 1065
MEIER, F. C., see GILBERT, W. W.
MEINECKE, E. P., Experiments with repeating pine rusts, 327–342

Melanconium sacchari, sugar-cane rind rot, 647

MELHUS, I. E., see PORTER, D. R.; and W. J. HENDERSON, The yellow dwarf of onions, 86; C. S. REDDY, W. J. HENDERSON, and EDGAR VESTAL, A new virus disease, epidemic on onion, 73–77

Melons, rot due to Phytophthora, 85

Mercuric, chloride, acidulated, for control, potato Rhizoctonia, 713; chloride, ethyl, as a grain disinfectant, 80; compounds, toxicity in seed treatment and spraying, 86

Mercury, organic, dust for oat-smut control, 173

Method of testing the keeping quality of certain small fruits, 593–596

Mexico, tobacco-mosaic control in, 880

Microscopical technique, handbook, review of, 609

Migration of Bacterium tabacum through leaf tissues, 97

Mildew, downy, alfalfa, 651; of cotton, 648; powdery, barley, 651

MILLER, P. A., A disease of Lippia caused by Sclerotium rolfsii Sacc., 509–510; Progress report on studies of fire blight of apple, 86–97

Millet, further experiments with smut of, 413

MILLS, P. J., see EDGERTON, C. W.

MILLS, W. D., see THOMAS, H. E.

Minutes of the field meeting of the southern section of the American Phytopathological Society, July 11–13, 1928, 531

MIX, A. J., Further studies of privet anthracnose, 102; Further studies on Exoascaceae, 90

Moisture, effect of, on spore formation in **Corticium praticola**, sp. nov., 1069

Mold, black, of onions, 413

Monosporidial cultures of Ustilago zeae, infection by, 420–423; pathogenicity, 91

MONTSERIN, B. G., and J. G. COULSON, Acid sterilization of liquid culture media, 413

MORITA, HELENE T., see GODFREY, G. H.

Morphology, **Phoma terrestris**, n. sp., 699

Mosaic, cucumber, transmission by peach aphid, 109; diseases of plants, 413; local symptoms in leaves of Nicotiana, 92–93; masking of, in sweet potato, 933–942; of cucumber, 85; of Easter lily, 311–315; reaction of Chinese cucumbers to, 85–86; sugar cane, 646; symptoms on raspberry, 89; tobacco, control in Mexico, 880; tobacco, transmission by peach aphid, 109; tomato plants, separation of toxins from, 108; tomato, purification of virus of, 108; virus of tobacco, freeing from accompanying solids, 107–108

MOUNCE, IRENE, The biology of Fomes pinicola (Sw.) Cooke, 414

Multisporidial cultures of Ustilago zeae, pathogenicity of, 91

MUNCIE, J. H., and M. K. PATEL, Potency and specificity of a lytic principle (bacteriophage) obtained from Pseudomonas tumefaciens, 98

Musa, host of Pestalotia leprogena, 217, 227

Mushroom root rot, see Armillaria mellea, 869

Mutation, in Gloeosporium, 605; in Helminthosporium sativum, 155; in Ustilago zeae, 106; modification of sexual reaction in Ustilago zeae, 431–433; of Fusarium spp., 143

Mutations in **Corticium praticola**, sp. nov., 1088–1090

Mycelium, diploid, of Ustilago zeae, 430–431

Mycosphaerella brassicicola (Duby) Lindau, ring spot disease of cabbage, 972

Myrica, asplenifolia, host of Cronartium comptoniae, 453; californicula, significance in distribution of rust, 465; gale, not associated in western United States with Pinus ponderosa, 463–464

Myzus persicae, as an agent in virus transmission, 109

Narcissus, relation of hot-water treatment to basal rot of bulbs, 100

NEBLETTE, C. B., see TAUBENHAUS, J. J.

Necrosis, net, of potato tubers, 82

Nectria, cinnabarina, a twig canker of apple caused by, 1125–1128; **N. tropica** (Wr.), comb. nov., on coffee root, 970

NELSON, RALPH M., and J. A. BEAL, Experiments with bluestain fungi in southern pines, 1101–1106
Nematode, description of **Tylenchus brachyurus**, n. sp., 617; root knot, effects of some environmental factors on, 83–84; root lesion on pineapple, 94; **Tylenchus brachyurus**, n. sp., relation to host tissues, 622
Nematospora, coryli, 479–480; coryli, occurrence and insect transmission of, in California, 1138; on pomegranates, citrus, and cotton in California, 479–482
Nerium, host of Pestalotia versicolor, 223, 227
Net blotch, barley, 650
Net necrosis of potato tubers, 82
New Sclerospora from Fiji, 961–967
NEWHALL, A. G., and J. D. WILSON, A preliminary report on forced-air ventilation for the control of Cladosporium leaf mold of greenhouse tomatoes, 83
Nicotiana, spp., local symptoms of mosaic in leaves of, 92–93; tabacum, migration of Bacterium tabacum through the leaf tissues of, 97
NISHIDA, TŌJI, biography of, 881–883; in memoriam, 527
Nitrogen, organic, effect of, on growth of **Corticium praticola**, sp. nov., 1068–1069
NOLLA, J. A. B., Biologic control of the aphids Rhopalosihum persicae Sulzer and Aphis gossypii Glover, 102; The black shank of tobacco in Porto Rico, 93–94
Notes, Phytopathological, 687–689; plant-disease, from Central Andes, 969–974
NUCKOLS, S. B., and C. M. TOMPKINS, An undescribed leaf condition associated with damping-off diseases of sugar-beet seedlings, 317–318; Delayed thinning as a means of controlling the damping off of sugar beets, 95
Nuclei, behavior in Sorosporium reilianum, 435–436; behavior in Ustilago zeae, 419–420, 424–428

Oak, Armillaria mellea on, 872; host of Pestalotia glandicola, 206; parasitism of Conopholus on roots of, 102
Oat, smut control, 102–103; smuts, dusting seed for control of, 173
Oats, Fusarium head blight of, 143; Pythium spp., infecting roots, 552; relative resistance to smut, 413
Observations on Armillaria root rot of orchard trees, 869–873
O'DONNELL, FRANK GETCHELL, in memoriam, 527
Officers and representatives of the American Phytopathological Society, 512
OGILVIE, LAWRENCE, and CARL E. F. GUTERMAN, A mosaic disease of the Easter lily, 311–315
Oleander, host of Pestalotia versicolor, 223
Onion, a new virus disease, epidemic on, 73–77; black mold, 733; pink-root disease, 233–268; relation to H-ion concentration, 233–268; yellow dwarf, 86
Onions, black mold of, 413; **Phoma terrestris**, n. sp., as the cause of pink root of, 1135–1136; pink-root disease, 691
Ophiobolus graminis, chemical and morphological phenomena attending infection by, 103–104; Sacc., histological studies of wheat infected with, 414; soil phases in relation to, 103
Orchard trees, Armillaria root rot of, 869–873
Ovulariopsis gossypii, cotton mildew, 648
OWENS, CHARLES ELMER, Principles of plant pathology, book review, 177
Oxalis stricta L., new host of sugar-beet curly top, 1031
Oxygen, effect of, on spore formation in **Corticium praticola**, sp. nov., 1073–1074
Ozonium omnivorum, see Phymatotrichum omnivorum.

Pacific division of the American Phytopathological Society, report of the thirteenth annual meeting of, 1135–1147
Palms, host of Pestalotia palmarum, 210–212
Parasite, root-hair, of pineapple, 367–381
Parasites, bacterial, influence of environment on, 96
Parasitism, Cronartium comptoniae, 742; of Conopholis americana on oak roots, 102

PARSONS, H. E., see LEACH, J. G.
PATEL, M. K., see MUNCIE, J. H. Some researches on Pseudomonas tumefaciens, 98–99; Viability of certain plant pathogenes in soils, 295–300
Pathogenicity and cultural behavior of Ustilago zeae (Bekm.) Ung. from different localities, 885–916; of multisporidial and monosporidial cultures of Ustilago zeae, 91
Pathological aspects of Godronia cassandrae Peck, 1017
PATTERSON, Mrs. FLORA WAMBAUGH, in memoriam, 528
PAXTON, G. E., see SIDERIS, C. P.
Peach, aphid (Myzus persicae Sulz.) as an agent in virus transmission, 109–123; bacterial spot, control of, 106–107; phony disease, 107; wound gum of, 413
Peltandra, host of Pestalotia aquatica, 224, 227
Penicillia of Citrus, further studies on, 1144
Penicillium, digitatum, hydrogen-ion-concentration studies on, 1142; digitatum, on citrus fruits in transit, 89–90; italicum, hydrogen-ion-concentration studies on, 1142; italicum, on citrus fruits in transit, 89–90; oxalicum, on corn, 105
Peronospora trifoliorum, alfalfa downy mildew, 651
Peru, diseases, economic plants in, 645–656; stem rust of wheat in, 1041
Pestolotia, glandicola, 206, 227; host index, 227; key to species, 225–227; monograph of genus, Part I, 191–232; rhododendri, 215, 227; sphaerelloides, 206, 227; vaccinii, 201, 227
Pestalozzia, see monograph of Pestalotia (synonym), 191–232; synonym of Pestalotia, 193
PETRE, A. W., see VINSON, C. G.
Petroleum oils, saturated, as insecticides on living plants, some limiting factors in the use of, 1136
Phacidium infestans on conifers, 91
Phaseolus, limensis Macf., host, **Phoma terrestris**, n. sp., 703; vulgaris, effects of grafting on resistance and susceptibility to Colletotrichum lindemuthianum, 875–877
Phases, certain developmental, common to many fungi, 1117–1123
Phoenix, host of Pestalotia palmarum, 210, 227
Phoma abietena, 984; pitya, 983–984; **P. terrestris**, n. sp., distribution, 701; **P. terrestris**, n. sp., as the cause of pink root of onions, 1135–1136; **P. terrestris**, n. sp., pathogene, pink-root disease, onions, 699
Phomopsis, abietena, 987; citri, on citrus fruits in transit, 89–90; on lemon bark, discovery of the perfect stage of, 1138; pseudotsugae, history and distribution, 979–992
Phony disease of peach, 107
Phyllachora gratissima Rehm on avocado, 972
Phyllosticta narcissi, 1138
Phymatotrichum omnivorum, 1025; omnivorum, method of inoculation for, 167; root-rot investigations, a method of inoculation for, 167–170
Physiologic, forms, of Helminthosporium sativum, mutation in, 155; forms of Puccinia graminis in relation to its epidemiology, 951–959; specialization, in Fusarium spp., causing head blight of small grains, 143–154
Physiological, strains of Alternaria solani, 533–548; studies of Bacterium translucens, 99
Physiology and cytology of Ustilago zeae and Sorosporium reilianum, studies in the, 415–442; of Ustilago zeae and Sorosporium reilianum, 91
Phytopathogenic bacteria, agglutination tests with, 99
Phytopathological Society, see American Phytopathological Society
Phytopathology, 493–494, 504–505
Phytophthora, causing pink rot of potato, 92; infestans, 651; nicotianae, on tobacco, 93–94; rot of honeydew melons, due to a species of, 85
PIERCE, LESLIÉ, see ROBERTS, JOHN W.

PIERCE, W. H., and C. W. HUNGERFORD, A note on the longevity of the bean-mosaic virus, 605–606

PIERSTORFF, A. L., and J. D. SAYRE, Further results of oat-smut control in Ohio, 102–103

Pine, needles, brown-spot disease of, 993–999; repeating rusts of, 327–342; western white, canker, 575; white, needle blight of, 413; Woodgate rust of, 414

Pineapple, disease of, 367–381; nematode root lesion, 94; root disease of, due to **Tylenchus brachyurus**, n. sp., 611–629; root-observing boxes for, 407–412; root rots caused by species of Fusarium, 1146; stem rot of, 1146

Pineapples, pythiaceous root parasites of, 1145–1146

Pines, southern, experiments with bluestain fungi in, 1101–1106

Pink, root disease, onions, 691; root of onions, **Phoma terrestris**, n. sp., as the cause of, 1135–1136; rot of potato caused by Phytophthora, 92

Pinus, contorta, attacked by Cronartium comptoniae, 453–465; echinulata, as host for bluestain fungi, 1103; monticola, Dasyscypha fusco-sanguinea infecting, 575; ponderosa, not associated in western United States with Myrica gale, 463–464; rigida, as host for bluestain fungi, 1103; strobus, as host of Dasyscypha agassizii, 277; strobus, blister rust, 269–283; sylvestris, infection by Cronartium comptoniae, 741

Plant, disease notes from the Central Andes II, 969–974; quarantine procedure, 495–497; quarantines and the State, 487–492

Planting stock and disease-free seed, 500

Plants, economic, diseases of, in Peru, 645–656

Plasmodiophora vascularum, spore development of, 91–92

Pleosphaerulina briosiana, alfalfa leaf spot, 651

Plot, arrangement to distribute neighbor influence in field trials, 565–574; technique, 565

Podocarpus, host of Pestalotia cesatii, 202, 227

Pokka bong, of sugar cane, 343–366; susceptibility of sugar-cane varieties to, 343–366

Polymorphism of Leptosphaeria coniothyrium (Fckl.) Sacc., 879

Polyporus, balsameus Pk., cause of butt-rot, balsam fir, 725; circinatus Fr., cause of root and butt-rot of conifers, 745

Pomegranates, Nematospora on, 479–482

POOLE, R. F., Further studies on the control of stem rot of sweet potatoes, 84

PORTER, D. R., and I. E. MELHUS, Further studies on watermelon wilt in Iowa, 84

PORTER, R. H., Reaction of Chinese cucumbers to mosaic, 85–86; T. F. YU, and H. K. CHEN, Response of hulless barley to seed treatment for covered smut and stripe disease, 657–666

Potato, abnormality in, 175; apical leaf roll of, 82–83; early blight, 652; late blight, 651; pink rot of, caused by Phytophthora, 92; powdery scab, 651; relation of Bordeaux to the internal leaf temperature of, 943–949; spindle tuber, 1045; spindle-tuber disease of, 1137; tuber rots, 652; tuber transmission of psyllid yellows of, 1140; tubers, net necrosis of, 82; tubers, strains of Alternaria solani from, 533; virus diseases, 652; wart, 652; wilt, 652

Potatoes, sweet, control of stem rot of, 84

Potency of lytic principle from Pseudomonas tumefaciens, 98

Powdery mildew of raspberry in California, observations on, 1144–1145

"Pressure death point," various organisms, 750

Prevention of stem rust of wheat, sulphur dusting for, 631–643

Principles of plant pathology, book review, 177

PRIODE, C. N., Pokkah-bong and twisted-top diseases of sugar cane, 343–366

Privet, anthracnose, 102

Problems, present-day, 493–501

Prune, Armillaria mellea on, 872; diamond canker of, 1139

Pseudofumago, as a developmental phase of fungi, 1117–1123; phase of Leptosphaeria coniothyrium, 879
Pseudomonas, beticolum, viability in soils, 295–300; citriputealis, viability in soils, 295–300; marginatum, viability in soils, 295–300; phaseoli, viability in soils, 295–300; tabacum, viability in soils, 295–300; tumefaciens, see Bacterium tumefaciens; tumefaciens, viability in soils, 295–300
Pseudopeziza medicaginis, alfalfa leaf spot, 651
Pseudosaccharomyces, as a developmental phase of fungi, 1117–1123; phase of Leptosphaeria coniothyrium, 879
Pseudotsuga, taxicolia, Armillaria mellea on, 869; taxifolia (Lam.) Britt., canker on, 979
Psyllid yellows, tuber transmission of, in California, 1140
Puccinia, graminis, heterothallism in, 1146–1147; graminis Pers., physiological forms in relation to epidemiology, 951–959; graminis, relation of weather to development of, 1–71; graminis, stem rust, 649; graminis, stem rust, barley, 650; graminis tritici in Peru, 1041; iridis, on Iris, 101; purpurea, varietal susceptibility of sorghum varieties to, 104; sorghi, corn rust, 649; triticina, wheat leaf rust, 650
Purification of tomato-mosaic virus, 108
Pustule, bacterial, of soy bean, 96
Pyrheliometer used with beet leaves, 474
Pyrus malus, fire-blight control and Bordeaux spray, 285–293
Pythiaceae, taxonomic studies in the family, 1142–1143
Pythiaceous root parasites, of pineapples, 1145–1146; of various agricultural plants, 1140
Pythium spp., sugar cane, root rot and, 549

Quarantine procedure, plant, 495–497
Quarantines, plant, 487–492
Quercus, bicolor, parasitism of Conopholus americana on roots of, 102; garryana, Armillaria mellea on, 872; host of Pestalotia glandicola, 206, 227

Radiobacter group of bacteria, differentiation of the crown-gall type of bacteria from the nonpathogenic bacteria of, 98
Radish, bacterial spot of, 97
RAMSEY, G. B., and ALICE ALLEN BAILEY, The development of soil rot of tomatoes during transit and marketing, 383–390
Raspberries, spur blight of, 413
Raspberry, anthracnose of, a new, 601; in California, observations on powdery mildew of, 1144–1145; mosaic, 89
RAWLINS, T. E., see JONES, WALTER
REA, H. E., see TAUBENHAUS, J. J.
Red, clover, black-stem disease, 507–509; rot of sugar cane, 646, 971–972
Reduction during spore germination of Ustilago zeae, 424–428
Relation, moisture to stomatal infection, 751
Report, cotton-root-rot conference, College Station, Texas, 687–689; of action of council, 525–526; of advertising manager of Phytopathology, 517–518; of advisory board of American Phytopathological Society, 518–519; of business manager, 1928, of Phytopathology, 514–516; of committee on editing phytopathological abstracts, 520–521; of committee on endowment for Phytopathology, 519–520; of committee on foreign pests and plant diseases, 521–523; of editor in chief of Phytopathology, 516–517; of New York meeting of American Phytopathological Society, 511–512; of other business, 526; of representative on council of the A. A. A. S., 524–525; of resolutions committee, 524–525; of secretary-treasurer, 1928, of American Phytopathological Society, 512–514; of the thirteenth annual meeting of the Pacific Division of the American Phytopathological Society, 1135–1147; of the twentieth anniversary meeting of the American Phytopathological Society, 511–526; on phytopathological classics, 523–524
Resistance, and susceptibility of beans to Colletotrichum lindemuthianum, 875–877; of corn to cold, 105–106; of oat varieties to smut, 413; relative, of varieties and species of Citrus to bark dis-

eases, 1136; to Armillaria mellea, studies on the nature of, 1140–1141; to barley scald, inheritance of, 1141; to mosaic, breeding for, in cucumbers, 85; to Septoria tritici in wheat, 1139–1140
Response of hulless barley to seed treatment for covered smut and stripe disease, 657–666
Retinospora, host of Pestalotia funerea, 203, 227
Rhizidiocystis ananasi Sideris, geographical distribution of, 367–381; of pineapples, 367–381
Rhizoctonia, **Corticium praticola,** n. sp., study of the biology of a new spore-forming, 1059–1099; potatoes, control by acidulated mercuric chloride, 713; solani, control, 587; solani Kühn, control of, 383–389; solani Kühn, distribution of, 383–389; solani Kühn, of tomatoes, 383–389; solani, Lobelia disease due to, 585
Rhizoctonosis of Lobelia, 585–588
Rhododendron, host of Pestalotia macrotricha, 214–215, 227; host of Pestalotia rhododendri, 216, 227
Rhopalosihum persicae, biologic control of, 102
Rhynchosporium secalis, 1141; barley leaf scald, 651; inoculation and culture studies, 104
Ribes, death by white-pine-blister rust a phase of control, 274; relation to white-pine-blister rust, 269–283; scale for measuring, 399–405
Rice, Ustilaginoidea infecting, occurrence in United States, 591
RICHARDS, B. L., White spot of alfalfa and its relation to irrigation, 125–141
RIKER, A. J., see WRIGHT, W. H.; Studies on the influence of environment on infection by certain bacterial plant parasites, 96; G. W. KEITT, and W. M. BANFIELD, A progress report on the control of crown gall, hairy root, and other malformations at the unions of grafted apple trees, 483–486; W. M. BANFIELD, and G. W. KEITT, The relation of certain bacteria to the development of roots, 107
RIKER, REGINA S., see JONES, L. R.
Rind rot of sugar cane, 647
Ring spot, of cabbage, 972; of sugar cane, 646
ROBERTS, JOHN W., and LESLIE PIERCE, A promising spray for the control of peach bacterial spot, 106–107
Robinia pseudacacia, brooming disease of, 163–166
RODENHISER, H. A., see CHRISTENSEN, J. J.
Roguing, spindle tuber in seed-potato plats, 1045
Root, and butt-rot of conifers caused by Polyporus circinatus Fr., 745–746; boxes for observing, 407–412; disease, coffee, 653, 969; disease of pineapple and other plants due to **Tylenchus brachyurus,** n. sp., 611–629; disease of sugar cane, Pythium spp. and, 549; disease, sugar cane, 647; knot, cotton, 649; -knot nematode, effects of some environmental factors on, 83–84; susceptibility of cowpea varieties and selections to, 1145; lesion, nematode, 94; -observation boxes, 407–412; parasites, pythiaceous, of pineapples, 1145–1146; parasites, pythiaceous, of various agricultural plants, 1140; rot, cotton, report conference College Station, Texas, 687–689; rot of cotton, study of, 1025–1029; rot, Phymatotrichum, 167; rots, corn, 649; rots, pineapple, caused by species of Fusarium, 1146
Roots, relation of certain bacteria to development of, 107
Rosa sp., Leptosphaeria coniothyrium on, 879
Rose bay, host of Pestalotia macrotricha, 214–215; host of Pestalotia rhododendri, 216
Rosellinia root disease, coffee, 653
Roses, aerial "crown gall" of, 1145
Rot, corn-cob, 549
Rot, cotton-root, report conference College Station, Texas, 687–689; Leptosphaeria foot, of wheat in Alberta, 689–690; of honeydew melons, 85; of narcissus bulbs, 99–100; of narcissus bulbs, relation to hot-water treatment, 100; of potato, caused by Phytophthora, 92; root, Phymatotrichum, 167
Rots, potato-tuber, 652; seedling and root of corn, 649

RUDOLPH, B. A., Preliminary report on a digest of the world's literature on hadromycosis due to Verticillium spp., 1138
RUMBOLD, CAROLINE T., Blue-staining fungi found in the United States, 597–599
RUSSELL, R. C., Histological studies of wheat infected with Ophiobolus graminis Sacc., 414
Rust, alfalfa, 651; corn, 649; diseases of apple, 87; of cereals, relation of weather to development of, 1–71; of Iris, 101; of sorghum, 104; stem, barley, 650; stem, of wheat, 649; stem, of wheat, sulphur dusting to prevent, 631–643; stem, relation of physiologic forms to epidemiology of, 951–959; stem, resistant segregates from wheat crosses between two susceptible parents, 1129–1130; wheat leaf, 650; Woodgate, 414

SAGEN, H. E., see WRIGHT, W. H.
Saltation in Helminthosporium sativum, 155
Salts, inorganic, effect of, on growth of **Corticium praticola**, sp. nov., 1067–1068
SAMSON, R. W., see KRAYBILL, H. R.
SAMUELS, CHAS. D., see KNIGHT, HUGH
Sassafras, host of Pestalotia gracilis, 217, 227
SAYRE, J. D., see PIERSTORFF, A. L.
Scab, apple, relation of fall applications of fungicides to control, 87–88; Fusarium, of wheat and oats, 143; of cereals, development in 1928, 108; powdery, potato, 651; studies of fungicidal action of dusts and sprays in control on apple, 87
Scald, barley leaf, 651; of cereals caused by Rhynchosporium secalis, 104
Scale for measuring leaves, 399–405
SCHNEIDERHAN, F. J., The influence of the form and proportion of lime and copper sulphate on the suspension of Bordeaux mixture, 88
SCHULTZ, E. S., and REINER BONDE, Apical leaf roll of potato, 82–83
Scirpus, host of Pestalotia scirpina, 209, 227
Sclerophoma magnusiana, 987
Sclerospora, graminicola, hosts of, 391–397; graminicola, occurrence of, in Wisconsin, 391–397; new, from Australia, 1107–1115; **S. noblei**, n. sp., characteristics of, 1109; **S. noblei**, n. sp., diagnosis of, 1112–1113; **S. noblei**, n. sp., effect of, on host, 1107; **S. noblei**, n. sp., identity of, 1111; **S. northi**, n. sp., cause of leaf-splitting disease on Erianthus maximus in Fiji, 961–967
Sclerotial fungi, cultures of, 176
Sclerotinia, cultures of, 176; intermedia, compared with minor in culture, 301–309; minor, compared with intermedia in culture, 301–309; sclerotium, in connection with the green rot of apricots, 1136–1137
Sclerotium rolfsii on Lippia, 509–510
Second decade of the American Phytopathological Society, 503–506
Seed, grain disinfectant, ethyl-mercury chloride as, 80; infection by Bacterium phaseoli, 96; oats, dusting for smut control, 173; potato plots, roguing for spindle tuber, 1045; treatment, response of hulless barley to, for covered smut and stripe disease, 657–666; treatment with mercury and copper compounds, 86
Seedling, and root rots, corn, 649; blight, influence of environment on predisposition of wheat and corn strains to, 79; blight of corn, inheritance of resistance to, 79–80; disease of tobacco, caused by Septomyxa affinis, 90
Segregation during spore germination of Ustilago zeae, 424–428
SELLSCHOP, JACQ. P. F., A mutation in Glocosporium, 605
Separation of toxins from mosaic-tomato plants, 108
Septomyxa affinis on tobacco, 90
Septoria, acicola, brown-spot disease, pine needles, 993–999; acicola, control of, 998; acicola, host range, distribution, 997; acicola, isolation of, 994; tritici, resistance to, in wheat, 1139–1140
SEVERIN, HENRY H. P., see SWEZY, OLIVE
Sexual reaction, modification of, in Ustilago zeae, 431–433
SHAPOVALOV, MICHAEL, Tuber transmission of psyllid yellows in California, 1140

SHEAR, C. L., Life history of Sphaceloma ampelinum, 673–679; see FAWCETT, H. S.; and HENRY F. BAIN, Life history and pathological aspects of Godronia cassandrae Peck (Fusicoccum putrefaciens Shear) on cranberry, 1017–1024
Sheath disease, sugar-cane, 647
SHERBAKOFF, C. D., Preface to Leonian paper on Fusarium, 753; Verticillium wilt of cotton, 94
SIDERIS, C. P., Effect of the H-ion concentration of culture solution on behavior of Fusarium cromyophthoron and Allium cepa and development of pink-root disease symptoms, 233–268; Pineapple root rots caused by species of Fusarium, 1146; Pythiaceous root parasites of various agricultural plants, 1140; **Rhizidiocystis ananasi** Sideris, nov. gen. et sp., a root-hair parasite of pineapples, 367–382; Stem rot of pineapple plants, 1146; Taxonomic studies in the family Pythiaceae, 1142–1143; and G. E. PAXTON, Pythiaceous root parasites of pineapples, 1145–1146
Single spores, method of isolation from individual basidia of **Corticium praticola**, sp. nov., 1078–1084
Smilax, host of Pestalotia clavata, 220, 227
SMITH, C. O., Leaf and scape spot of the Amarillidaceae in California, 1138–1139
SMITH, RALPH E., Diamond canker of the prune, 1139; Life history of Sclerotinia sclerotiorum in connection with the green rot of apricots, 1136–1137
Smut, corn, 649; covered, and stripe disease, response of hulless barley to seed treatment for, 657–666; covered, control on barley, 81; covered, wheat, 650; false, of maize, 589; loose, barley, 651; loose, of wheat, 103, 650; millet, 413; of corn, 415–452; of oat, control, 102–103; pathogenicity and cultural characteristics of Ustilago zeae, 885–916; relation of removal of smut balls to control in winter wheat, 79; relative resistance of oat varieties to, 413; stinking, control on wheat, 81
SNELL, WALTER H., Some observations upon the white-pine-blister rust in New York, 269–283
Snow cover, a disease of conifers related to, 91
Snowberry, anthracnose of, 101–102
Soil phases of wheat take-all problem, 103
Soils, viability of pathogenes in, 295–300
Some present-day problems, 493–501
Sore shin of cotton, 648
Sorghum, bacterial leaf spot of, 82; plumosum, host for **Sclerospora noblei**, n. sp., 1107–1115; Pythium spp., infecting roots, 552; varietal susceptibility to Puccinia purpurea, 104
Sorosporium reilianum, 415–442; physiology and cytology of, 91; studies in physiology and cytology of, 413
Soy bean, bacterial pustule of, 96
Specialization, physiologic, in Fusarium spp., 143
Specificity of lytic principle from Pseudomonas tumefaciens, 98
Sphaceloma ampelinum, life history of, 673–679
Spindle tuber, of the potato, 1137; relation to roguing in seed-potato plats, 1045
Spongospora subterranea, 651
Spore germination, of Sorosporium reilianum, 435–436; of Ustilago zeae, 419–420; of Ustilago zeae, reduction and segregation during, 424–428
Spores of Plasmodiophora vascularum, development of, 91–92
Sporidia of Ustilago zeae from infected corn leaves, 433–434
Sporophores, Polyporus circinatus Fr., 747
Sporotrichum, fruit spot and surface rot of apple, 443–452; malorum, isolation of, 445; malorum, morphology of, 445–446; malorum, pathogenicity of, 447–448; malorum, symptoms on apple produced by, 443–444
Spot, bacterial, of radish and turnip, 97; brown, coffee fruit and leaf, 653; eye, coffee, 652; fruit, tomato caused by Aplanobacter michiganense, 690; leaf, sycamore, caused by Stigmina platani, 667–671

SPRAGUE, RODERICK, Host-range and life-history studies of some leguminous Ascochytae, 917–932
Spray for control of peach bacterial spot, 106–107
Spraying, calcium monosulphide a substitute for lime sulphur, 106
Sprays, effect on weight of cherry fruit, 88; fungicidal action in apple-scab control, 87
Spread of Cronartium comptoniae, 462–464
STAKMAN, E. C., see LAMBERT, E. B.; J. J. CHRISTENSEN, and W. F. HANNA, Mutation in Ustilago zeae, 106; M. N. LEVINE, and J. M. WALLACE, The value of physiologic-form surveys in the study of the epidemiology of stem rust, 951–959
STARRETT, RUTH COLVIN, A new host of sugar-beet curly top, 1031–1035
Stem, rot, of pineapple plants, 1146; rot, of sweet potatoes, studies on control of, 84; rust, barley, 650; rust, relation of weather to development, 1–71; rust, wheat, 649; rust, wheat, in Peru, 1041
Stemphylium, leaf spot of tomatoes, 92
Sterilization of liquid culture media, 413
STEVENS, N. E., A method of testing the keeping quality of certain small fruits, 593–596; and HENRY F. BAIN, Storage rots of cranberries in the 1928 crop, 1037–1039
STEWART, GEORGE, Stem-rust-resistant segregates from wheat crosses between two susceptible parents, 1129–1130
Stigmina platani, leaf spot of sycamore caused by, 667–671
Stilbella flavida, coffee eye spot, 652
STILLINGER, C. R., Dasyscypha fusco-sanguinea Rehm on western white pine, Pinus monticola Dougl., 575–584
Stinking smut, control by removal of smut balls, 79
Stomatal infection, relation to moisture, 751
STONE, R. E., Further experiments with millet smut, 413; Observations on the relative resistance to smut infection of some varieties of oats, 413; see HOWITT, J. E.
Storage rots, cranberries, 1037
Stripe disease, control of, on barley, 81; response of hulless barley to seed treatment for covered smut and, 657–666
Studies, in the physiology and cytology of Ustilago zeae and Sorosporium reilianum, 415–442; on the mycelium of Cronartium comptoniae Arthur on Pinus sylvestris L., 741–744; on the nature of host resistance to Armillaria mellea, 1140–1141; on the variability and dissociations in the genus Fusarium, 753–868
Substitute for lime sulphur, calcium monosulphide as, 106
Sugar beet, attenuation of curly-top virus, 975–977; curly top, new host of, 1031; curly-top virus, modification of, by its various hosts, 1141–1142; dry-rot canker, 94–95; infected with curly top, filterable microorganism from, 1143; leaf condition with damping off, 317–318; thinning as a means of controlling damping off, 95
Sugar beets, ecological studies of curly top, 467–477; mass action in relation to infection with special reference to curly top of, 1137
Sugar cane, a new genus of Dematiaceae on, 1051; chloroses, 647; diseases of, 343–366; eye spot, 646; mosaic, 646; red rot of, 971–972; rind rot, 647; ring spot, 646; root disease, 647; root rot and Pythium spp., 544; sheath disease, 647; top rot and red rot of, 971–972; susceptibility of varieties of, 343–366
Sugars as food for **Corticium praticola**, sp. nov., 1067
Sulphur, chemistry of toxic factor of, 89; dusting for prevention of stem rust of wheat, 631–643; mixtures, adhesiveness of, 89
Surface rot of apple, caused by Sporotrichum fruit spot, 443–452
Susceptibility, and resistance of beans to Colletotrichum lindemuthianum, 875–877; of corn to cold, 105–106; of cowpea varieties and selections to Fusarium

wilt and root knot, 1145; of varieties of sorghum to Puccinia purpurea, 104
Sweet, clover, black-stem disease of, 507–509; potato, masking of mosaic symptoms, 933–942; potato, soil rot or pox caused by Actinomyces, undescribed species, 179–190; potatoes, control of stem rot of, 84
SWEZY, OLIVE, and HENRY H. P. SEVERIN, A filterable microorganism from Eutettix tenellus and the sugar beet, both infected with curly top, 1143
Sycamore, leaf spot, caused by Stigmina platani, 667–671
Symphoricarpos albus, anthracnose of, 101–102
Symptoms of mosaic on species of Nicotiana, 92–93
Synandrospadix, host of Pestalotia pallidicolor, 217, 227
Synchytrium endobioticum, 652

Take-all of wheat, relation of soil phases, 103
TAPKE, V. F., The rôle of humidity in the life cycle, distribution, and control of the loose-smut fungus of wheat, 103
TAUBENHAUS, J. J., B. F. DANA, W. N. EZEKIEL, W. J. BACH, and J. P. LUSK, A method of inoculation for Phymatotrichum root-rot investigations, 167–170; W. N. EZEKIEL, and C. B. NEBLETTE, Airplane photography in the study of cotton root rot, 1025–1029; W. N. EZEKIEL, and H. E. REA, A new cotton wilt, 171–173
Taxonomic studies in the family Pythiaceae, 1142–1143
Taxonomy, Polyporus balsameus Pk., 728; Septoria acicola, 995
Tea, host of Pestalotia theae, 213
Temperature, effect of, on growth of **Corticium praticola,** sp. nov., 1070; effect on physiologic forms of Fusarium spp., 143; influence on mutation in Helminthosporium sativum, 155; of potato leaflets as influenced by Bordeaux mixture, 943–949; relations, cranberry end rot, 1023; relations, **Phoma terrestris,** n. sp., 697; small fruit keeping quality affected by, 593
TENG, S. C., Rhizoctonosis of Lobelia, 585–588
Thea, host of Pestalotia theae, 213, 227
Thielaviopsis paradoxa (Des.) v. Holn., on sugar cane, 972
THOMAS, HAROLD E., Studies on the nature of host resistance to Armillaria mellea, 1140–1141; and A. B. BURRELL, A twig canker of apple caused by Nectria cinnabarina, 1125–1128; and W. D. MILLS, Rust diseases of apple, 87
THOMAS, RALPH CLEON, biography of, 1057–1058; in memoriam, 529
Thuja, host of Pestalotia funerea, 203, 227
TILFORD, PAUL E., and CURTIS MAY, The effect of Bordeaux mixture on the internal temperature of potato leaflets, 943–949
Tilletia, caries, effect of, on development of wheat ear, 681–685; laevis, covered smut of wheat, 650; tritici (Bjerk.) Wint., effect of, on development of wheat ear, 681–685
Timothy, reaction in agronomic strains to Ustilago striaeformis, 105
TIMS, E. C., see EDGERTON, C. W.
TISDALE, L. E., see YOUNG, H. C.
TISDALE, W. B., A disease of tobacco seedlings caused by Septomyxa affinis (Sherb.) Wr., 90; Progress in the control of black shank of tobacco through disease resistance, 93
TISDALE, W. H., and W. N. CANNON, Ethylmercury chloride as a seed-grain disinfectant, 80
Tobacco, angular leaf spot, 93; blackfire, 93; black shank, control through disease resistance, 93; black shank in Porto Rico, 93–94; mosaic control in Mexico, 880; mosaic virus, freeing from accompanying solids, 107–108; mosaic virus, inactivated by high pressures, 749; seedling disease, 90; transmission of mosaic virus by peach aphid, 109
Tomato, disease of, 383–389; diseases in Peru, 655; fruit spot, caused by Aplanobacter michiganense, 690; leaf mold,

973; leaf mold, control by forced-air ventilation, 83; mosaic plants, separation of toxins from, 108; purification of virus of mosaic of, 108; Stemphylium leaf spot, 92
TOMPKINS, C. M., Further studies on the dry-rot canker of sugar beets, 94–95; see NUCKOLS, S. B.
Top rot of sugar cane, 971
TORO, RAFAEL A., Plant-disease notes from the Central Andes II, 969–974
Toxic factor of sulphur, 89
Toxicity, of mercury and copper compounds, 86; of some commercial dusts to conidia of Bremia lactucae, 1143–1144
Toxins which produce some typical mosaic symptoms, separation from mosaic-tomato plants, 108
Transmission, insect, of Nematospora in California, 1138; tuber, of psyllid yellows in California, 1140
Transportation and market pathology, 500
Treatment, seed, response of hulless barley to, for covered smut and stripe disease, 657–666
TU, CHIH, Physiologic specialization in Fusarium spp. causing head blight of small grains, 143–154; see CHRISTENSEN, J. J.
Tumor on potato stem, 175
Turnip, bacterial spot of, 97
Twisted top, cause of, 343–366; of sugar cane, 343–366
Tylenchus brachyurus, n. sp., diagnosis, 618; **brachyurus**, n. sp., environmental relations of, 626; **brachyurus**, n. sp., root disease of pineapples due to, 611–629; T. sp., cause of root lesion on pineapple, 94

Uromyces medicaginis, alfalfa rust, 651
Ustilaginoidea virens (Cke.) Tak., maize false smut due to, 589
Ustilago, nuda, loose smut, barley, 651; striaeformis, reaction in agronomic strains of timothy to, 105; tritici, loose smut, wheat, 650; tritici, relation of humidity to, 103; zeae, 415–442; zeae (Bekm.) Ung., pathogenicity and cultural behavior, 885–916; zeae, corn smut, 649; zeae, mutation in, 106; zeae, pathogenicity of multisporidial and monosporidial cultures, 91; zeae, physiology and cytology of, 91; zeae, studies in physiology and cytology of, 413

Vaccinium, host of Pestalotia vaccinii, 201, 227
Vacuolar changes in diseased cells, 95
VALLEAU, W. D., Are black fire and angular leaf spot of tobacco identical? 93; Tobacco-mosaic control in Mexico, 880; and E. N. FERGUS, Black-stem disease of alfalfa, sweet clover, and red clover, 507–509
Value of physiologic-form surveys in the study of the epidemiology of stem rust, 951–959
Variability and dissociations in the genus Fusarium, 753–868
Variation in cultures of Alternaria solani, 533
Vector, insect, of aster-yellows virus, 1009–1015
Ventilation, forced-air, for control of tomato-leaf mold, 83
Venturia inaequalis, studies of fungicidal action of dusts and sprays in control of, 87
Verticillium, alboatrum, on cotton, 94; spp., preliminary report of a digest of the world's literature on hadromycosis due to, 1138; wilt, of cotton, 94
Vigna sinensis, host, **Phoma terrestris**, n. sp., 703
VINSON, C. G., and A. W. PETRE, Progress in freeing the virus of mosaic disease of tobacco from accompanying solids, 107–108
Virus, aster-yellows, 1009–1015; disease, on onion, 73–77; diseases, potato, 652; of tobacco mosaic, freeing from solids, 107–108; of tomato mosaic, purification, 108; sugar-beet, attenuation of, by resistant varieties, 975–977; transmission by peach aphid, 109
Vitis, host of Pestalotia menezesiana, 219, 227; host of Pestalotia pezizoides, 197–227

WAGENER, W. W., Lentinus lepideus Fr., cause of heart rot of living pines, 705–712
WALLACE, J. M., see STAKMAN, E. C.
Walnut, Armillaria mellea on, 872
Wart, potato, 652
Watermelon, host of Pestalotia torulosa, 222; wilt, 84
Watermelons, chemical injury to, 85
Weather, relation to development of stem rust, 1–71
WEBER, GEORGE F., A Stemphylium leaf spot of tomatoes, 92
Weight of cherry fruit, effect of sprays on, 88
WEISS, FREEMAN, The basal rot of narcissus bulbs caused by Fusarium sp., 99–100; The relation of the hot-water treatment of narcissus bulbs to basal rot, 100
WERNER, H. O., Relation of time of roguing to spread of spindle tuber in seed-potato plats, 1045–1049
Western white pine, canker, 575
WESTON, W. A. R. DILLON, Effect of Tilletia caries (DC.) Tul. (T. tritici (Bjerk.) Wint.) on development of wheat ear, 681–685
WESTON, WILLIAM H., JR., A new Sclerospora from Fiji, 961–967; A new Sclerospora from Australia, 1107–1115; The occurrence of Sclerospora graminicola on maize in Wisconsin, 391–397
Wheat, chemical and morphological phenomena attending infection by Ophiobolus graminis, 103–104; control of stinking smut in, 79, 81; covered smut, 650; crosses between two susceptible parents, stem-rust-resistant segregates from, 1129–1130; effect of Tilletia caries on development of ear, 681–685; Fusarium head blight of, 143; infection of, with Ophiobolus graminis Sacc., 414; leaf rust, 650; Leptosphaeria foot rot of, in Alberta, 689–690; loose smut, 650; loose-smut fungus, relation of humidity to, 103; Pythium spp. infecting roots, 552; resistance to Septoria tritici in, 1139–1140; seedling blight, 79; stem rust, 649; stem rust in Peru, 1041; stem rust, physiologic-form surveys in the study of epidemiology of, 951–959; sulphur dusting for prevention stem rust of, 631–643; take-all, relation of soil phases, 103
WHETZEL, H. H., Cultures of sclerotial fungi, 176; S. E. A. MCCALLAN, and T. C. LOH, Calcium arsenate as a fungicide, 83
WHITE, HAROLD E., and M. W. GARDNER, Bacterial spot of radish and turnip, 97
White, pine, as host of Dasyscypha agassizii, 277; -pine blister rust, 269–283; spot, of alfalfa, 125
WHITNEY, W. A., see HARTER, L. L.
WILLIAMS, R. C., and H. C. YOUNG, Chemistry of the toxic factor of sulphur, 89
WILLISON, R. S., The relation of wound gum to fungous invasion in peach and grape, 413
Willow, new disease, in Canada and New England, 413
WILSON, E. E., see KEITT, G. W.; and G. W. KEITT, The effect of sprays on the weight of cherry fruit, 88
WILSON, J. D., see NEWHALL, A. G.
WILSON, MALCOLM, and GLENN GARDNER HAHN, The history and distribution of Phomopsis pseudotsugae in Europe, 979–992
Wilt, bacterial, of Cruciferae, 413; coffee, 653; cotton, 94, 647; of aster, control, 101; of asters, 100–101; of cotton, a new, 171; of watermelon, 84; potato, 652; -resistant asters, 100–101
WINGARD, S. A., biography of Ralph Cleon Thomas, 1057–1058
Wire-screen fences for control of aster yellows, 100
Witches' broom on black locust, 163
Wither tip of mango, 973
Wood, blue staining of, fungi causing, 597
Woodgate rust, distribution of, in Canada, 414
WRIGHT, WILLIAM H., A. J. RIKER, and H. E. SAGEN, Studies on the differentiation of the crown-gall type of bacteria from nonpathogenic bacteria of the radiobacter group, 98; and W. H. BANFIELD,

Studies on the bacteriological differentiation of the crown-gall and hairy-root types of bacteria, 97–98

Yellow dwarf of onions, 86
Yellows, of aster, control, 101; of aster, control by wire-screen fences, 100
YOUNG, H. C., Dusts composed of lime sulphur and sulphur, 88–89; see WILLIAMS, R. C.; and L. E. TISDALE, Adhesiveness of sulphur mixtures, 89
YOUNG, V. H., see HIGGINS, B. B.
YU, T. F., see PORTER, R. H.

ZAUMEYER, W. J., Seed infection by Bacterium phaseoli, 96
Zea mays, Ustilaginoidea virens infecting, 589
ZELLER, S. M., Another anthracnose of raspberry, Gloeosporium, 601–603; see CHILDS, LEROY
Zinc sulphate for control of peach bacterial spot, 106–107